

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.</p>					
1. REPORT DATE (DD-MM-YYYY) 10/13/2017		2. REPORT TYPE Abstract		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE A Rare Case of Periapillary Tumor				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Capt Peter C Johnson				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg 4430 JBASA-Lackland, TX 78236-9908 210-292-7141				8. PERFORMING ORGANIZATION REPORT NUMBER 17311	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg 4430 JBASA-Lackland, TX 78236-9908 210-292-7141				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release. Distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Title: A Rare Case of Periapillary Tumor Peter Johnson ¹ , MD, Ryan Cho ² , MD, Kelly A. Haeusler ³ , MD, Katherine M. Cebe ³ , MD, Angelo H. Paredes ² , MD. 1. Department of Internal Medicine, San Antonio Military Medical Center, San Antonio Uniformed Services Health Education Consortium, Fort Sam Houston, TX 2. Department of Internal Medicine, Gastroenterology Service, San Antonio Uniformed Services Health Education Consortium, Fort Sam Houston, TX 3. Department of Pathology, San Antonio Uniformed Services Health Education Consortium, Fort Sam Houston, TX Introduction: Gangliocytic paraganglioma is a rare tumor that can affect anywhere in the gastrointestinal tract, though are most frequently found in the second portion of the duodenum. They commonly present with abdominal pain, gastrointestinal bleeding, or rarely with obstructive jaundice.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Clarice Longoria
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code) 210-292-7141

Title: A Rare Case of Periapillary Tumor

Peter Johnson¹, MD, Ryan Cho², MD, Kelly A. Haeusler³, MD, Katherine M. Cebe³, MD, Angelo H. Paredes², MD.

1. Department of Internal Medicine, San Antonio Military Medical Center, San Antonio Uniformed Services Health Education Consortium, Fort Sam Houston, TX
2. Department of Internal Medicine, Gastroenterology Service, San Antonio Uniformed Services Health Education Consortium, Fort Sam Houston, TX
3. Department of Pathology, San Antonio Uniformed Services Health Education Consortium, Fort Sam Houston, TX

Introduction: Gangliocytic paraganglioma is a rare tumor that can affect anywhere in the gastrointestinal tract, though are most frequently found in the second portion of the duodenum. They commonly present with abdominal pain, gastrointestinal bleeding, or rarely with obstructive jaundice. This is a case of an asymptomatic periapillary gangliocytic paraganglioma.

Case: A 45year old male with a history gastroesophageal reflux well controlled on a PPI was referred iron deficiency anemia with no overt bleeding. He underwent an upper endoscopy that revealed a submucosal nodule involving the ampulla (Figure 1). The endoscopic ultrasound subsequently revealed a 2.3x1.5cm hypoechoic heterogeneous mass of the periapillary region (Figure 2). Fine needle aspiration showed a neoplasm with neuroendocrine differentiation. The immunohistochemical stain was strongly positive for synaptophysin and negative chromogranin, S-100, CD-117, and DOG-1. The periapillary mass was resected using endoscopic mucosal resection, with the final pathology consistent with gangliocytic paraganglioma (Figure 3). The patient had further testing with negative urine metanephrines and catecholamines. There was no radiographic evidence of distal metastatic disease.

Discussion: Gangliocytic paraganglioma (GP) is a rare neuroendocrine tumor, with only around 200 cases reported. It consisted on three distinct cellular elements: spindle cells, epithelial cells, and ganglion cells. In spindle-cell predominant tumors, the differential diagnosis includes schwannomas and GIST. When visualized endoscopically, GPs are difficult to distinguish from other submucosal tumors and may appear as a pedunculated or sessile polypoid mass that may be smooth or ulcerated. Endoscopic biopsies are frequently negative since the tumor is submucosal. An EUS is frequently necessary to obtain a fine-needle aspiration as well as evaluate for local spread of the tumor and detect metastatic lymph nodes. The majority of GPs are benign without local or distant spread. However, lymph nodes metastases and distant metastases to other organs have been reported. Treatment is dependent on the size of the tumor and the presence of any metastatic lymph nodes or other metastatic sites. Case reports have shown that overall the prognosis is good but there have been reported deaths from metastatic tumors. In conclusion, we present the case of an incidental periapillary gangliocytic paraganglioma.

*The views expressed are those of the author/presenter and do not reflect the official views or policy of the Department of Defense or its components.

FIGURE 1:



Figure 1: Endoscopic image of the periampullary mass

FIGURE 2:



Figure 2: Endoscopic ultrasound image of the periampullary mass

FIGURE 3:




Figure 3: Seven ganglion cells have abundant cytoplasm, round nuclei, and prominent nucleoli (small arrows). Spindle cells are arranged in fascicles (large arrows). Epithelioid endocrine cells are arranged in trabeculae (arrowheads)